

EUROPHYSICS LETTERS

Europhys. Lett., (), pp. ()

Dephasing in a quantum dot due to coupling with a quantum point contact

YEHOSHUA LEVINSON

Weizmann Institute of Science, 76100 Rehovot, Israel

(received ; accepted in final form)

PACS. 73.23.-b – Mesoscopic systems.

PACS. 72.10.-d – Theory of electronic transport; scattering mechanisms.

Abstract. – We consider the dephasing of an one-electron state in a quantum dot due to charge fluctuations in a biased quantum point contact coupled to the dot capacitively. The contribution to the dephasing rate due to the bias depends on temperature and bias in the same way as shot-noise in the point contact at zero frequency, but do not follow the $|t|^2(1 - |t|^2)$ suppression.

Recently, it became possible to measure the phase and hence also the coherence of resonance electron transmission through a (pinched) quantum dot (QD) [1, 2, 3, 4]. Experiments of this type are performed using Aharonov-Bohm rings loaded by a QD in one of its arms, the ring being imbedded in a two- or four-terminal configuration. The conductance of such a system contains a term oscillating with magnetic flux Φ threaded through the ring and this term is sensitive to the phase of the transmission amplitude t_{QD} through the QD.

If the transmission through the QD is coherent one can calculate $\mathcal{G}$ from the Landauer formula in terms of the transmission through the ring $\mathcal{T}$. The transmission amplitude t_{QD} through a pinched QD is small (it is true even for resonance transmission, since generically the QD is asymmetric) and for small t_{QD} one finds $|\mathcal{T}|^2 = |\mathcal{T}|_0^2 + \text{Re}\{a^* t_{QD} \exp(2\pi i \Phi / \Phi_0)\}$, where Φ_0 is the flux quanta. The two terms correspond to the non-oscillating and oscillating parts of $\mathcal{G}$ in the two-terminal configuration. The first is the transmission through the "free" arm and the second is due to the interference between the (multiple) electron paths in the "free" and "loaded" arms. The effective "amplitude" a is responsible for the geometry of the ring outside the QD and of the external contacts to the ring. When the temperature $T = 0$ all energy depending transmission amplitudes has to be calculated at the Fermi energy ϵ_F and for $T \neq 0$ one has to average over thermal energies near ϵ_F . The phase of transmittance through a QD in an Aharonov-Bohm ring was analyzed in papers [5, 6] and also in connection with the "phase slippage" [7].

When the coherence of the electron states in the QD is destructed due to interaction with some "environment" the amplitude t_{QD} has to be replaced by its average $\langle t_{QD} \rangle$ with respect to states of the "environment". The destruction of coherence is not necessary related to inelastic scattering [8] and this is why we call it "dephasing".

Typeset using EURO-L^AT_EX

One source of such a dephasing can be the capacitive coupling between the QD and a quantum point contact (PC) which is close in space to the QD [9, 10]. Charge fluctuations in the PC create a fluctuating potential at the QD and modulate the electron states in the QD. This modulation being random in time dephases the states in the QD and destructs the coherence of the transmission through the QD.

The charge fluctuations in the PC depend on whether the PC is in equilibrium or supports a current due to an applied bias V . Hence one can expect that the visibility of the ring conductance oscillations with flux will change with current in the PC.

The aim of this work is to calculate the dephasing time of a state in a QD induced by its capacitive coupling to a PC. We will be interested mainly in the additional contribution to the dephasing rate which is due to the current in the PC. To simplify the problem we consider dephasing in an isolated QD not coupled to leads and coupled capacitively to a PC.

To make things more transparent assume only one state in the QD with energy ϵ_0 and only one channel in the point contact with coordinate x along the channel and an effective one dimensional barrier potential $U(x)$. For simplicity we assume the PC to be symmetric with respect to $x = 0$ (the maximum of the barrier).

The Hamiltonian of the QD is $H_{QD} = \epsilon_0 c^+ c$, where c is an operator removing one electron from the QD state.

The Hamiltonian of the PC is ($\hbar = 1$)

$$H_{PC} = \int dx \psi^+(x) \left[-\frac{1}{2m} \frac{d^2}{dx^2} + U(x) \right] \psi(x) = \sum_k \epsilon_k [a_k^+ a_k + b_k^+ b_k]. \quad (1)$$

Here $\psi(x)$ is the electron field operator, which can be presented as [11]

$$\psi(x) = \sum_k [a_k \chi_k(x) + b_k \phi_k(x)], \quad (2)$$

where $\chi_k(x)$ and $\phi_k(x)$ are scattering states radiated from the left and right ohmic contacts a and b and belonging to the same energy ϵ_k and a_k, b_k are the corresponding electron operators. The scattering states have the following properties ($k > 0$)

$$\begin{aligned} \chi_k(x) &= L^{-1/2} \{ \exp(+ikx) + r_k \exp(-ikx) \}, & \text{for } x \rightarrow -\infty, \\ \chi_k(x) &= L^{-1/2} t_k \exp(+ikx), & \text{for } x \rightarrow +\infty, \end{aligned} \quad (3)$$

r_k and t_k being the reflection and transmission coefficients defined by the potential barrier $U(x)$ and L is some normalization length. Similar equations can be written for $\phi(x) = \chi(-x)$.

The Hamiltonian of the interaction between the QD and PC, assumed to be weak, is

$$H_{int} = c^+ c W \equiv c^+ c \int dx \delta U(x) \psi^+(x) \psi(x). \quad (4)$$

If one combines $H_{PC} + H_{int}$ one can see that $\delta U(x)$ is the average change of the barrier potential $U(x)$ due to one electron occupying the QD state when $\langle c^+ c \rangle = 1$. This change is localized near the squeezing of the PC $x = 0$. If one combines $H_{QD} + H_{int}$ one can see that W is the change of the QD state energy ϵ_0 due to the interaction of the electron in the QD with the electron density in the PC.

The dephasing of the QD state is described by the relaxation of the average state amplitude $\langle c(t) \rangle$, where $c(t) = \exp(+iHt) c \exp(-iHt)$ is the amplitude in Heisenberg representation with the total Hamiltonian $H = H_{QD} + H_{PC} + H_{int}$.

The Heisenberg amplitude equation of motion is

$$(d/dt)c(t) = i \exp(+iHt) [H, c] \exp(-iHt) = -i[\epsilon_0 + W(t)]c(t), \quad (5)$$

with

$$W(t) = \exp(+iHt)W \exp(-iHt). \quad (6)$$

Eq.(5) demonstrates that $W(t)$ is indeed the time depending modulation of the energy level ϵ_0 . Solving this equation and averaging one finds

$$\langle c(t) \rangle = \left\langle c(0) \exp(-i\epsilon_0 t) T_t \exp \left\{ -i \int_0^t dt W(t) \right\} \right\rangle, \quad (7)$$

where T_t means time ordering. We decouple the average in Eq.(7) approximating the time ordered exponent by a Gaussian exponent and obtain

$$\langle c(t) \rangle = \langle c(0) \rangle \exp(-i\epsilon_0 t) \exp(-\Phi(t)), \quad (8)$$

where

$$\Phi(t) = \frac{1}{2} \int_0^t dt' \int_0^t dt'' K(t' - t''), \quad (9)$$

and $K(t)$ is the quantum correlator of the level modulation

$$K(t) = \frac{1}{2} [\langle W(t)W(0) \rangle + \langle W(0)W(t) \rangle]. \quad (10)$$

We also approximate $W(t)$ defined in Eq.(6) by

$$W(t) = \exp(+iH_{PC}t)W \exp(-iH_{PC}t) \quad (11)$$

neglecting the influence of the QD on the fluctuations in the PC. With this approximation the average in Eq.(10) reduces to the average with respect to the state of the PC (in equilibrium or with current) which plays now the role of an environment. This last approximation means that we have in mind an experimental situation when the current in the QD used to measure its linear conductance $\mathcal{G}$ is small while the current in the PC can be relatively large.

The correlator $K(t)$ decays in time with some time scale τ_c which is the correlation time of the QD state energy modulation. For large $t \gg \tau_c$ one find from Eq.(8)

$$\langle c(t) \rangle = \langle c(0) \rangle \exp(-i\epsilon_0 t) \exp(-t/\tau_\varphi) \quad (12)$$

with

$$\frac{1}{\tau_\varphi} = \frac{1}{2} \int_{-\infty}^{+\infty} dt K(t). \quad (13)$$

Hence the *average amplitude* of the QD state decays. Note that under same conditions the population of the QD state is constant $\langle c^\dagger(t)c(t) \rangle = \langle c^\dagger(0)c(0) \rangle$, since H_{int} commutes with the number of electrons in the QD $c^\dagger c$. Same is true for the energy of the QD $\epsilon_0 c^\dagger c$. This means that the decay of $\langle c(t) \rangle$ with the time constant τ_φ is indeed *dephasing* and not *energy relaxation* or *escape* from the QD.

Using Eq.(4) and Eq.(2) one finds the energy modulation in terms of scattering states

$$W = \sum_{k,k'} [a_k^\dagger a_{k'} A_{k,k'} + b_k^\dagger b_{k'} B_{k,k'} + a_k^\dagger b_{k'} C_{k,k'} + b_k^\dagger a_{k'} \bar{C}_{k,k'}] \quad (14)$$

with integrals describing the interaction between the QD and the PC

$$\begin{aligned} A_{k,k'} &= \int dx \delta U(x) \chi_k(x)^* \chi_{k'}(x), & B_{k,k'} &= \int dx \delta U(x) \phi_k(x)^* \phi_{k'}(x), \\ C_{k,k'} &= \int dx \delta U(x) \chi_k(x)^* \phi_{k'}(x), & \bar{C}_{k,k'} &= \int dx \delta U(x) \phi_k(x)^* \chi_{k'}(x) = C_{k',k}^*. \end{aligned} \quad (15)$$

Using Eq.(14) one can find (similar to the calculation [12] of the current correlator in a PC)

$$\begin{aligned} < W(t)W > = \sum_{k,k'} \exp[i(\epsilon_k - \epsilon_{k'})t] \times \\ & \{n_k^a(1 - n_{k'}^a)|A_{k,k'}|^2 + n_k^b(1 - n_{k'}^b)|B_{k,k'}|^2 + n_k^a(1 - n_{k'}^b)|C_{k,k'}|^2 + n_k^b(1 - n_{k'}^a)|\bar{C}_{k,k'}|^2\}, \end{aligned} \quad (16)$$

n_k^a and n_k^b being the occupation numbers for states with energy ϵ_k in the left and in the right ohmic contacts of the PC a and b respectively,

$$n_k^{a,b} = [\exp[-(\epsilon_k - \mu_{a,b})/T] + 1]^{-1}, \quad (17)$$

where $\mu_{a,b}$ are the electrochemical potentials in the ohmic contacts a, b and $V = -(\mu_a - \mu_b)/e$ is the applied voltage between a and b (assuming $e > 0$).

The spectral density of the QD level modulation can be calculated using Eq.(10) and Eq.(16) as follows

$$\begin{aligned} S(\omega) &= \frac{1}{2\pi} \int_{-\infty}^{+\infty} dt \exp(i\omega t) K(t) = \frac{1}{2} \sum_{k,k'} \delta(\epsilon_k - \epsilon_{k'} + \omega) \times \\ & \{n_k^a(1 - n_{k'}^a)|A_{k,k'}|^2 + n_k^b(1 - n_{k'}^b)|B_{k,k'}|^2 + n_k^a(1 - n_{k'}^b)|C_{k,k'}|^2 + n_k^b(1 - n_{k'}^a)|\bar{C}_{k,k'}|^2\} + \\ & \{\omega \rightarrow -\omega\}. \end{aligned} \quad (18)$$

To simplify the spectral density assume that the squeezing in the PC (i.e. the barrier $U(x)$) is described by only one length scale d , the energy spacing between the thresholds for different channels in the PC being $\Delta\epsilon \simeq 1/md^2 \ll \epsilon_F$. With this assumption the scale in k for the coefficients A, B, C and $\bar{C}$ is d^{-1} . Assume also that the bias in the PC and the temperature are small in the sense that $eV, T \ll \Delta\epsilon$. One can see that with these assumptions the relevant frequencies ω are much smaller than the inverse of the time of flight through the PC and that the main contribution come from states close to the Fermi level. Since $|\epsilon_k - \epsilon_{k'}| \simeq v_F |k - k'| \simeq \omega$ (where v_F is the Fermi velocity) one finds that for $\omega \ll v_F/d \ll \epsilon_F$ the relevant $|k - k'| \ll d^{-1}$ and one can replace in the coefficients A, B, C and $\bar{C}$ the momenta k and k' by k_F . After this replacement we just skip the momenta in the notation and note that $|C|^2 = |\bar{C}|^2$.

Changing summation over $k > 0$ to integration over ϵ_k and introducing a function [13]

$$F(\omega) = \int \int d\epsilon d\epsilon' [\delta(\epsilon - \epsilon' + \omega) + \delta(\epsilon - \epsilon' - \omega)] n_\epsilon (1 - n_{\epsilon'}) = F(-\omega) \quad (19)$$

one finds the explicit expression for the spectral density

$$S(\omega) = \frac{L^2}{8\pi^2 v_F^2} [(|A|^2 + |B|^2)F(\omega) + |C|^2(F(\omega + eV) + F(\omega - eV))], \quad (20)$$

Subtracting the equilibrium part of the spectral density (corresponding to $V = 0$) we find the spectral density of the QD energy modulation due to the current

$$S_V(\omega) = \lambda [F(\omega + eV) + F(\omega - eV) - 2F(\omega)], \quad (21)$$

where we defined a coupling constant $\lambda = (L^2/8\pi^2 v_F^2)|C|^2$.

According to Eq.(21) the spectra of the QD energy modulation due to nonequilibrium charge fluctuations in the PC is the same as the shot-noise spectra in the PC [12, 13]. The typical modulation frequencies are determined by the energy window near ϵ_F where the electron fluxes from the left and right ohmic contacts a and b of the PC do not compensate, i.e. $\omega \simeq eV/\hbar$ for high bias $eV \gg T$ and $\omega \simeq T/\hbar$ for low bias $eV \ll T$. One can check that for low enough temperature and bias $T, eV \ll \Delta\epsilon$ these frequencies are indeed smaller than the inverse time of flight. These frequencies also define the correlation time of the QD level modulation $\tau_c \simeq \omega^{-1}$.

However the amplitude of the energy modulation determined by the coupling constant λ depends on the transmittance through the PC in a way very different from that of the shot noise, which is proportional to $|t|^2(1 - |t|^2)$. To see it we can rewrite the coupling constant in terms of changes in r and t introduced by the perturbation $\delta U(x)$ to the potential $U(x)$. Using the representation of the Green function for the Schrödinger equation with potential $U(x)$ in terms of scattering states

$$G(x, x') = (mL/ikt)\chi(x_>)\phi(x_<), \quad (22)$$

where $x_>$ and $x_<$ are the larger and the smaller of x and x' , and also the properties of the scattering states (for a symmetric barrier)

$$\chi^*(x) = r^*\chi(x) + t^*\phi(x), \quad (23)$$

one can find

$$\lambda = (8\pi^2)^{-1}|r^*\delta t + t^*\delta r_a|^2 = (8\pi^2)^{-1}|r^*\delta t + t^*\delta r_b|^2, \quad (24)$$

where δr_a and δr_b are the changes of r for waves coming from ohmic contacts a and b and t is the change in t .

Let see how the coupling constant depends on the barrier in the PC. If the barrier $U(x)$ is infinite ($|t|^2 = 0$) the functions χ and ϕ do not overlap and it follows from Eq.(15) that $|C|^2 = 0$ and $\lambda = 0$. Same can be obtained from Eq.(24) since zero transmittance of an infinite barrier can not be changed by a final perturbation and hence from $t = 0$ it follows that $\delta t = 0$. It means that what matters for the nonequilibrium dephasing in the QD is not the applied bias in the PC, but the current. In case of an infinite barrier applied bias does not change the electron density and its fluctuations in the ohmic contacts a and b .

If there is no barrier ($|t|^2 = 1$), i.e. the quantum point contact is replaced by a quantum wire, χ and ϕ are plane waves and one finds from Eq.(24) and Eq.(15)

$$\lambda = \frac{1}{8\pi^2}|\delta r_a|^2 = \frac{1}{8\pi^2}|\delta r_b|^2 = \frac{1}{8\pi^2 v_F^2} \left| \int dx \delta U(x) \exp(i2k_F x) \right|^2. \quad (25)$$

In terms of the spectral density using Eq.(21) the nonequilibrium contribution to dephasing rate can be written as

$$\tau_\varphi^{-1} = \pi S_V(0) = 2\pi\lambda[F(eV) - F(0)]. \quad (26)$$

One can easily estimate that this contribution $\tau_\varphi^{-1} \simeq \lambda eV/\hbar$ for high bias $eV \gg T$ and $\tau_\varphi^{-1} \simeq \lambda(eV)^2/T\hbar$ for low bias $eV \ll T$. The dependence of the dephasing time on bias and temperature follows the shot-noise spectral intensity for zero frequency [12]. Note that $\hbar/\tau_\varphi$ is much smaller than the rms average modulation of the QD level $<(\delta\epsilon_0)^2>^{1/2} \simeq <W^2>^{1/2}$ because of the effect of dynamical narrowing, i.e. $<(\delta\epsilon_0)^2>^{1/2} \tau_c \ll 1$. The last inequality is valid because of $\lambda \ll 1$.

The limiting expressions for λ demonstrates that in spite of the fact that the dephasing is originated by the same charge fluctuations that are responsible for shot noise, the dephasing rate does not follow generally the intensity of shot noise which is proportional to $|t|^2(1 - |t|^2)$.

A similar calculation of the equilibrium part of modulation spectra results in

$$S_0(\omega) = 2(\mu + \lambda)F(\omega), \quad (27)$$

where

$$2\mu = (L^2/8\pi^2 v_F^2)[|A|^2 + |B|^2] = (8\pi^2)^{-1}[|r^* \delta r_a + t^* \delta t|^2 + |r^* \delta r_b + t^* \delta t|^2]. \quad (28)$$

A cutoff at the time-of-flight frequency v_F/d has to be introduced to this spectra, but the equilibrium contribution to the dephasing rate defined by $S_0(0)$ is independent on this cutoff and equal to $\tau_\varphi^{-1} = 2\pi(\mu + \lambda)T$. (Note that when the thermal current noise at zero frequency is calculated in this way it is in agreement with the d.c. conductance according to the Nyquist theorem).

If the QD is not pinched and the electron in state ϵ_0 has a finite escape rate to the leads Γ the dephasing rate competes with this rate. For zero dephasing the (nonaveraged) transmission amplitude t_{QD} contains a resonant factor $-i/[(\epsilon_F - \epsilon_0) + i\Gamma]$. Due to dephasing this factor is replaced in the averaged amplitude $\langle t_{QD} \rangle$ by

$$\int_0^\infty dt \exp[-\Gamma t - \Phi(t) + i(\epsilon_F - \epsilon_0)t]. \quad (29)$$

Generally the factor Eq.(29) defines a non Lorentzian line-shape but in the case of dynamical narrowing $\langle (\delta\epsilon_0)^2 \rangle^{1/2} \tau_c \ll 1$ it reduces to $-i/[(\epsilon_F - \epsilon_0) + i(\Gamma + \hbar/\tau_\varphi)]$ just adding the dephasing rate to the escape broadening.

Recently the problem of dephasing was also addressed in an unpublished paper of Aleiner, Wingreen and Meir [14] using a different approach.

This work was initiated as a result of discussions with M.Heiblum and E.Buks about the possibility of the so called "Which Path?" experiment in the Aharonov-Bohm interferometer. I acknowledge these discussions very much. I am thankful to Y.Imry for discussions related to level broadening and dephasing rate and also to A.Stern and F.von Oppen for discussing the results of this work.

REFERENCES

- [1] YACOBY A., HEIBLUM M., MAHALU D., and SHTRIKMAN H., *Phys.Rev.Lett*, **74** (1995) 4047.
- [2] YACOBY A., SHUSTER R., and HEIBLUM M., *Phys.Rev.B*, **53** (1996) 9583.
- [3] BUKS E., SHUSTER R., HEIBLUM M., MAHALU D., UMANSKY V., and SHTRIKMAN H., *Phys.Rev.Lett*, **77** (1996) 4664.
- [4] SHUSTER R., BUKS E., HEIBLUM M., MAHALU D., UMANSKY V., and SHTRIKMAN HADAS, *Nature*, **385** (1997) 417.
- [5] LEVY YEYATI A., and BUTTIKER M., *Phys.Rev.B*, **52** (1995) R14360.
- [6] HACKENBROICH G., and WEIDENMULLER H. A., *Phys.Rev.Lett.*, **76** (199) 110.
- [7] OREG Y., and GEFEN Y., *cond-mat/9607145*, (1996) .
- [8] STERN A., AHARONOV Y., and IMRY Y., *Phys.Rev.A*, **41** (1990) 3436.
- [9] FIELD M., SMITH C. G., PEPPER M., RITCHIE D. A., FROST J. E. F., JONES G. A. C., and HASKO D. G., *Phys.Rev.Lett*, **70** (1993) 1311.
- [10] HEIBLUM M., ET AL, *unpublished*.
- [11] BUTTIKER M., *Phys.Rev. B*, **46** (1992) 12485
- [12] LESOVIK G. B., *JETP Lett.*, **49** (1989) 592.
- [13] ERIC YANG S. R., *Solid State Comm.*, **81** (1992) 375.
- [14] ALEINER I. L., WINGREEN N. S., and MEIR Y., *cond-mat/9702001*, (1997) .